

# 114 Soybean

AMOUNT OF PRODUCT PER GALLON OF SPRAY MIXTURE					
GRAMINICIDES					
PRODUCT	RATE	ADDITIVES		COMMENTS	
		COC.	NIS		
ARROW <sup>1</sup>	0.65 oz +	1.3 oz	or	0.3 oz	<sup>1</sup> Label directions for application timing; restrictions for preharvest interval; and use of forage for spot treatments are the same as those for broadcast applications for ARROW, ASSURE II, FUSION, SECTION, SELECT, and SELECT MAX.  <sup>2</sup> Do not apply spot treatments of POAST or POAST PLUS in addition to broadcast treatments.
ASSURE II <sup>1</sup>	0.5 oz +	1.25 oz	or	0.3oz	
FUSION <sup>1</sup>	0.75 oz +	1.5 oz	or	0.5oz	
POAST <sup>2</sup>	1.9 oz +	1.3 oz		-----	
POAST PLUS <sup>2</sup>	2.9 oz +	1.3 oz		-----	
SECTION <sup>1</sup>	0.65 oz +	1.3 oz	or	0.3 oz	
SELECT <sup>1</sup>	0.65 oz +	1.3 oz	or	0.3 oz	
SELECT MAX <sup>1</sup>	0.85 oz +	1.3 oz	or	0.3 oz	
VOLUNTEER <sup>1</sup>	0.65 oz +	1.3 oz	or	0.3 oz	
<b>GLYPHOSATE</b>					
GLYPHOSATE FORMULATION <sup>3</sup>		RATE		COMMENTS	
3 lb ae /gal		0.64 to 2.56 oz		<sup>3</sup> Apply GLYPHOSATE products as a spot-treatment before initial pod set of soybean. Susceptible plants in the treated area may be killed or injured. Do not treat more than 10% of the total area to be harvested. Mix the low rate for annuals <6" tall and the high rate for certain hard-to-control perennials. Higher rates may be warranted for certain weeds. Consult label for rates for specific weeds. See Page 22 for list of glyphosate products.	
3.75 lb ae/gal		0.51 to 2.05 oz			
4 lb ae /gal		0.48 to 1.92 oz			
4.5 lb ae/gal		0.42 to 1.7 fl oz			
<b>PARAQUAT</b>					
PRODUCT	RATE	SURFACTANT		COMMENTS	
GRAMOXONE	0.33 to 0.67 oz	0.33 to 0.5 oz		Thoroughly wet foliage but not to the point of runoff. Do not use around home gardens, schools, recreational areas, golf courses, or playgrounds.	
FIRESTORM	0.33 to 0.67 oz	0.33 to 0.5 oz			
PARAZONE	0.33 to 0.67 oz	0.33 to 0.5 oz			

## PREHARVEST